

ANALYSIS OF UNBOUNDED OPERATORS AND RANDOM MOTION

PALLE E.T. JORGENSEN

UNIVERSITY OF IOWA

ABSTRACT. We study infinite weighted graphs with view to “limits at infinity,” or boundaries at infinity. Examples of such weighted graphs arise in infinite (in practice, that means “very” large) networks of resistors, or in statistical mechanics models for classical or quantum systems. But more generally our analysis includes reproducing kernel Hilbert spaces and associated operators on them. If X is some infinite set of vertices or nodes, in applications the essential ingredient going into the definition is a reproducing kernel Hilbert space; it measures the differences of functions on X evaluated on pairs of points in X . And the Hilbert norm-squared in $\mathcal{H}(X)$ will represent a suitable measure of energy. Associated unbounded operators will define a notion of dissipation, it can be a graph Laplacian, or a more abstract unbounded Hermitian operator defined from the reproducing kernel Hilbert space under study. We prove that there are two closed subspaces in reproducing kernel Hilbert space $\mathcal{H}(X)$ which measure quantitative notions of limits at infinity in X , one generalizes finite-energy harmonic functions in $\mathcal{H}(X)$, and the other a deficiency index of a natural operator in $\mathcal{H}(X)$ associated directly with the diffusion. We establish these results in the abstract, and we offer examples and applications. Our results are related to, but different from, potential theoretic notions of “boundaries” in more standard random walk models. Comparisons are made.

I. INTRODUCTION

We will use the theory of unbounded Hermitian operators with dense domain in Hilbert space in a study of infinite weighted graphs with view to “limits at infinity.” We begin with introducing the tools from

2000 *Mathematics Subject Classification.* Primary 47B25, 47B32, 47B37, 47S50, 60H25, 81S10, 81S25.

Key words and phrases. Operators in Hilbert space, deficiency spaces, networks of resistors, statistical mechanics models, harmonic functions, graphs, Schroedinger equations, selfadjoint extension operators, reproducing kernel Hilbert spaces.

Work supported in part by the US National Science Foundation.

operator theory as developed by H. M. Stone, John von Neumann, Kurt Friedrichs, and Tosio Kato with view to our particular setup. We stress that a Hermitian operator may not be selfadjoint, and that the discrepancy is measured by deficiency-indices (details below and the books [31] and [32], and more recently [11] and [28].) In physical problems, see e.g., [7], these mathematical notions of defect take the form of “boundary conditions;” for example waves that are diffracted on the boundary of a region in Euclidean space; the scattering of classical waves on a bounded obstacle [25]; a quantum mechanical “particle” in a repulsive potential that shoots to infinity in finite time; or in more recent applications (see e.g., [22], [8], [9], [27]) random walk on infinite weighted graphs G that “wander off” to points on an idealized boundary of G . In all of the instances, one is faced with a dynamical problem: For example, the solution to a Schrödinger equation, represents the time evolution of quantum states in a particular problem in atomic physics.

The operators in these applications will be Hermitian, but in order to solve the dynamical problems, one must first identify a self-adjoint extension of the initially given operator. Once that is done, von Neumann’s spectral theorem can then be applied to the selfadjoint operator. A choice of selfadjoint extension will have a spectral resolution, i.e., it is an integral of an orthogonal projection valued measure; with the different extensions representing different “physical” boundary conditions. Hence non-zero deficiency indices measure degrees of non-selfadjointness, and deficiency spaces “measure” boundary obstructions or scattering on an obstacle.

The variety of applied problems that lend themselves to computation of deficiency indices and the study of selfadjoint extensions are vast and diverse. As a result, it helps if one can identify additional structures that throw light on the problem.

Our results are inspired in part by the following recent developments in related areas: fractals in the small and in the large [13, 17, 19], representation theory [15, 16], operator algebras [10, 30], harmonic analysis [14, 17, 12, 20]; and multiresolutions/wavelets [1, 29, 23, 21].

We will further use notions from dynamics, infinite matrix products, to prove essential selfadjointness of families of Hermitian operators arising naturally in reproducing kernel Hilbert spaces. The latter include graph Laplacians for infinite weighted graphs (G, w) with the Laplacian in this context presented as a Hermitian operator in an associated Hilbert space of finite energy functions on the vertex set in G . Other examples include Hilbert spaces of band-limited signals.

Further applications enter into the techniques used in discrete simulations of stochastic integrals, see [18].

We encountered the present operator theoretic results in our study of discrete Laplacians, which in turn have part of its motivation in numerical analysis. A key tool in applying numerical analysis to solving partial differential equations is discretization, and use of repeated differences; see e.g., [6]. Specifically, one picks a grid size h , and then proceeds in steps: (1) Starting with a partial differential operator, then study an associated discretized operator with the use of repeated differences on the h -lattice in $\mathbb{R}^d$. (2) Solve the discretized problem for h fixed. (3) As h tends to zero, numerical analysts evaluate the resulting approximation limits, and they bound the error terms. For this purpose, one must use a metric, and the norm in Hilbert space has proved an effective tool, hence the Hilbert spaces and the operator theory.

This procedure connects to our present graph-Laplacians: When discretization is applied to the Laplace operator in d continuous variables, the result is the graph of integer points $\mathbb{Z}^d$ with constant weights. But if numerical analysis is applied instead to a continuous Laplace operator on a Riemannian manifold, the discretized Laplace operator will instead involve infinite graph with variable weights, so with vertices in other configurations than $\mathbb{Z}^d$.

Inside the technical sections we will use standard tools from analysis and probability. References to the fundamentals include [6], [24], [26] and [33].

There is a large literature covering the general theory of reproducing kernel Hilbert spaces and its applications, see e.g., [4], [2], [3], [5], and [34]. Such applications include potential theory, stochastic integration, and boundary value problems from PDEs among others. In brief summary, a reproducing kernel Hilbert space $\mathcal{H}$ consists of two things: a Hilbert space of functions f on a set X , and a reproducing kernel k , i.e., a complex valued function k on $X \times X$ such that for every x in X , the function $k(\cdot, x)$ is in $\mathcal{H}$ and reproduces the value $f(x)$ from the inner product $\langle k(\cdot, x), f \rangle$ in $\mathcal{H}$, so the formula

$$f(x) = \langle k(\cdot, x), f \rangle$$

holds for all x in X . Moreover, there is a set of axioms for a function k in two variables that characterizes precisely when it determines a reproducing kernel Hilbert space. And conversely there are necessary and sufficient conditions that apply to Hilbert spaces $\mathcal{H}$ and decide when $\mathcal{H}$ is a reproducing kernel Hilbert space.

Here we shall restrict these “reproducing” axioms and obtain instead a smaller class of reproducing kernel Hilbert spaces. We add two additional axioms: Firstly, we will be reproducing not the values themselves of the functions f in $\mathcal{H}$, but rather the differences $f(x) - f(y)$ for all

pairs of points in X ; and secondly we will impose one additional axiom to the effect that the Dirac mass at x is contained in $\mathcal{H}$ for all x in $\mathcal{H}$. When these two additional conditions are satisfied, we say that $\mathcal{H}$ is a relative reproducing kernel Hilbert space.

It is known that every weighted graph (the infinite case is of main concern here) induces a relative reproducing kernel Hilbert space, and an associated graph Laplacian. Under certain conditions, the converse holds as well: Given a relative reproducing kernel Hilbert space $\mathcal{H}$ on a set X , it is then possible in a canonical way to construct a weighted graph G such that X is the set of edges in G , and such that its energy Hilbert space coincides with $\mathcal{H}$ itself. In our construction, the surprise is that the edges in G as well as the weights on the edges may be built directly from only the Hilbert space axioms defining the initially given relative reproducing kernel Hilbert space. Since this includes all infinite graphs of electrical resistors and their potential theory (boundaries, harmonic functions, and graph Laplacians) the result has applications to these fields, and it serves to unify diverse branches in a vast research area.

II. OPERATOR THEORETIC FRAMEWORK

This section contains the precise definitions of the terms used above, and to be used in later sections: the particular discrete networks, the weights on edges, the associated Hilbert spaces, and the infinite Laplacians. We open with two lemmas which establish links between the graph theoretic networks on one side, and the operator theory (unbounded Hermitian operators) on the other. This will allow us to encode certain “boundaries” with two subspaces of an associated Hilbert space.

Let X be a set, and let $c : X \times X \rightarrow \mathbb{R}_{\geq 0}$ a function satisfying the following four conditions:

(i) For all $x \in X$,

$$\# \{y \in X | c(x, y) \neq 0\} < \infty.$$

(ii) Symmetry:

$$c(x, y) = c(y, x), \forall x, y \in X.$$

(iii)

$$c(x, x) = \sum_{y \in X} c(x, y).$$

- (iv) For all $x, y \in X$, such that $x \neq y$, there is a finite set $\{x_0, x_1, \dots, x_n\}$ of distinct points in X such that $c(x_i, x_{i+1}) \neq 0$, $0 \leq i < n$, and $x_0 = x$, $x_n = y$.

Lemma 2.1. *Let (X, c) be a system as described above with the function*

$$c : X \times X \rightarrow \mathbb{R}_{\geq 0}$$

satisfying (i)–(iv).

For $x \in X$, let δ_x be the Dirac-function on X supported at x .

Then there is a Hilbert space $\mathcal{H}(X, c)$ containing $\{\delta_x\}_{x \in X}$ with inner product $\langle \cdot, \cdot \rangle$ such that

$$\langle \delta_x, \delta_x \rangle = c(x, x), \quad (1)$$

and

$$\langle \delta_x, \delta_y \rangle = -c(x, y) \text{ if } x \neq y. \quad (2)$$

Proof. We will be working with functions on X modulo multiples of the constant function $\mathbb{1}$ on X . We define the Hilbert space $\mathcal{H}(X, c)$ as follows:

$$\mathcal{H}(X, c) := \{\text{all functions } f : X \rightarrow \mathbb{C} \text{ s.t.}$$

$$\sum_{\substack{x, y \\ \text{s.t. } x \neq y}} c(x, y) |f(x) - f(y)|^2 < \infty\}; \quad (3)$$

and we set

$$\begin{aligned} \|f\|^2 &= \langle f, f \rangle \\ &= \mathcal{E}(f) := \frac{1}{2} \sum_{x \neq y} \sum c(x, y) |f(x) - f(y)|^2 \text{ for } f \in \mathcal{H}(X, c). \end{aligned} \quad (4)$$

It is immediate that $\mathcal{H}(X, c)$ is then a Hilbert space, and that the Dirac-functions δ_x are in $\mathcal{H}(X, c)$. If $\langle \cdot, \cdot \rangle$ is the inner product corresponding to (4), then a direct computation shows that (1) and (2) are satisfied. $\square$

Lemma 2.2. *Let (X, c) be a system satisfying conditions (i)–(iv) above, and let $\mathcal{H}(X, c), \langle \cdot, \cdot \rangle$ be the Hilbert space introduced in Lemma 2.1.*

For elements $u \in \mathcal{H}(X, c)$, set

$$(\Delta u)(x) := \langle \delta_x, u \rangle \text{ for } x \in X. \quad (5)$$

Then Δ is a Hermitian operator with a dense domain $\mathcal{D}$ in $\mathcal{H} := \mathcal{H}(X, c)$.

Proof. Select some base-point o in X . Let $x \in X$, and select $(x_i)_{i=0}^n \subset X$ such that $x_{i-1} \neq x_i$, $c(x_{i-1}, x_i) \neq 0$, and $x_0 = o$, $x_n = x$. Then if $f \in \mathcal{H}$, we get the following estimate

$$|f(x) - f(o)| \leq \left(\sum_{i=1}^n \frac{1}{c(x_{i-1}, x_i)} \right)^{\frac{1}{2}} \|f\| \quad (6)$$

where $\|f\|$ is the norm in $\mathcal{H}$. By Riesz' lemma, there is a unique $v_x \in \mathcal{H}$ such that

$$f(x) - f(o) = \langle v_x, f \rangle \text{ for all } f \in \mathcal{H}. \quad (7)$$

Now let $\mathcal{D}$ be the linear span of $(v_x)_{x \in X}$. It follows from (7) that $\mathcal{D}$ is dense in $\mathcal{H}$, and that

$$\Delta v_x = \delta_x - \delta_o. \quad (8)$$

To prove (8) first note that the functions (v_x) in (7) must be real-valued. This is a consequence of the uniqueness in Riesz. We now verify (8) by the following computation:

$$\begin{aligned} (\Delta v_x)(y) &= (\text{by (5)}) \langle \delta_y, v_x \rangle \\ &= (\text{by (7)}) \delta_y(x) - \delta_y(o) \\ &= (\delta_x - \delta_o)(y), \end{aligned}$$

which is the desired conclusion (8).

A direct inspection yields the following two additional properties.

$$\begin{aligned} \langle \Delta u_1, u_2 \rangle &= \langle u_1, \Delta u_2 \rangle, \text{ and} \\ \langle u, \Delta u \rangle &\geq 0, \text{ for all } u_1, u_2, \text{ and } u \in \mathcal{D}. \end{aligned}$$

□

Definition 2.3.

(a) Let G^0 and G^1 be two sets (in our case the infinite cases are of main importance), and assume the following:

- $G^1 \subset G^0 \times G^0$.
- If $(x, y) \in G^1$, then $x \neq y$, and we write $x \sim y$.
- If $(x, y) \in G^1$ then $(y, x) \in G^1$.
- For every pair of points x and y in G^0 , there is a finite subset

$$\{(x_{i-1}, x_i) \mid i = 1, 2, \dots, n\} \subset G^1$$

such that $x_0 = x$ and $x_n = y$.

- For every $x \in G^0$ we assume that the set of neighbors is finite, i.e.,

$$\# \text{Nbh}_G(x) < \infty, \quad (9)$$

where

$$\# \text{Nbh}_G(x) = \{y \in G^0 | y \sim x\}. \quad (10)$$

- (b) Let $c: G^1 \rightarrow \mathbb{R}_+$ be a function satisfying

$$c(x, y) = c(y, x) \text{ for all } (x, y) \in G^1. \quad (11)$$

- (c) For functions $u: G^0 \rightarrow \mathbb{C}$ set

$$\mathcal{E}(u) := \frac{1}{2} \sum_{\substack{\text{all } xy \\ \text{such that } x \sim y}} c(x, y) |u(x) - u(y)|^2. \quad (12)$$

We will consider the Hilbert space $\mathcal{H}_E$ of all functions modulo constants such that

$$\mathcal{E}(u) < \infty; \quad (13)$$

called the *energy Hilbert space*, and (13) referring to *finite energy*.

For functions $u: G^0 \rightarrow \mathbb{C}$, we define the *graph Laplacian*, or simply the *Laplace operator* Δ by

$$(\Delta u)(x) = \sum_{y \sim x} c(x, y) (u(x) - u(y)). \quad (14)$$

Proposition 2.4.

- (a) The Laplace operator Δ in (14) is defined on a dense linear subspace $\mathcal{D}$ in $\mathcal{H}_E$, and Δ maps $\mathcal{D}$ into $\mathcal{H}_E$.
 (b) Pick some o in G^0 . Then for every $x \in G^0 \setminus \{o\}$, there is a unique $v_x \in \mathcal{H}_E$ such that

$$\langle v_x, u \rangle_E = u(x) - u(o) \text{ for all } u \in \mathcal{H}_E. \quad (15)$$

- (c) The function v_x in (15) satisfies

$$\Delta v_x = \delta_x - \delta_o. \quad (16)$$

- (d) For the subspace $\mathcal{D}$ in (a) we may take

$$\mathcal{D} = \text{span} \{v_x | x \in G^0 \setminus \{o\}\}, \quad (17)$$

i.e., all finite linear combinations of the v_x family of vectors.

(e) *The Dirac-functions δ_x in (16) satisfy*

$$\delta_x = c(x) v_x - \sum_{y \sim x} c(x, y) v_y \quad (18)$$

where

$$c(x) := \sum_{y \sim x} c(x, y). \quad (19)$$

(f) *The operator Δ is Hermitian on $\mathcal{D}$, i.e.,*

$$\langle \Delta u_1, u_2 \rangle = \langle u_1, \Delta u_2 \rangle \text{ holds for all } u_1, u_2 \in \mathcal{D}, \quad (20)$$

where

$$\begin{aligned} \langle u_1, u_2 \rangle &:= \langle u_1, u_2 \rangle_E \\ &:= \mathcal{E}(u_1, u_2) \\ &:= \frac{1}{2} \sum_{\substack{x \\ \text{such that } x \sim y}} \sum_y c(x, y) \left(\overline{u_1(x)} - \overline{u_1(y)} \right) (u_2(x) - u_2(y)). \end{aligned} \quad (21)$$

(g) *Semiboundedness:*

$$\langle u, \Delta u \rangle \geq 0 \text{ for all } u \in \mathcal{D}. \quad (22)$$

Proof.

(a) Follows from (b) and (c).

(b) This is an application of Riesz' lemma for the Hilbert space $\mathcal{H}_E$; see [22]. Indeed, pick $(x_i)_{i=0}^n \subset G^0$ such that $(x_{i-1}, x_i) \in G^1$, and $x_0 = 0$, $x_n = x$; then

$$|u(x) - u(o)| \leq \left(\sum_{i=1}^n \frac{1}{c(x_{i-1}, x_i)} \right)^{\frac{1}{2}} \|u\|_{\mathcal{H}_E} \quad (23)$$

where

$$\|u\|_{\mathcal{H}_E} := \mathcal{E}(u)^{\frac{1}{2}}, \quad u \in \mathcal{H}_E. \quad (24)$$

The desired conclusion (15) is then immediate from Riesz.

(c) It is clear from (15) that each function

$$v_x: G^0 \rightarrow \mathbb{R} \quad (25)$$

is real valued and that $\{v_x | x \in G^0 \setminus (0)\}$ span a dense subspace $\mathcal{D}$ in $\mathcal{H}_E$.

Hence to prove (16), we need to show that

$$\langle v_y, \Delta v_x - (\delta_x - \delta_0) \rangle = 0 \text{ for all } x, y \in G^0 \setminus (0). \quad (26)$$

Because of (12), we may impose the following re-normalization:

$$v_x(o) = 0, \quad \forall x \in G^0 \setminus (0). \quad (27)$$

With this, now (26) follows from (12), (14) and (15), by a direct computation which we leave to the reader; see also [22].

(d) Follows from (c).

(e) By (d) it is enough to prove that for all $z \in G^0$, we have the identity:

$$\left\langle v_z, \delta_x - c(x) v_x + \sum_{y \sim x} c(x, y) v_y \right\rangle = 0,$$

which in turn is a computation similar to the one used in (c).

(f) and (g) Follow by polarization, and a direct computation of (22). Indeed, if

$$u = \sum_x \xi_x v_x \quad (28)$$

is a finite summation over $x \in G^0 \setminus (0)$, $\xi_x \in \mathbb{C}$, then $\xi_x = (\Delta u)(x)$, and

$$\langle u, \Delta u \rangle = \sum_x |\Delta u(x)|^2 + \left| \sum_x (\Delta u)(x) \right|^2. \quad (29)$$

□

Definition 2.5. *The following two closed subspaces play a critical role in our understanding of geometric boundaries of weighted graphs (G, c) ; i.e., a graph with G^0 serving as the set of vertices, G^1 the set of edges, and*

$$c: G^1 \rightarrow \mathbb{R}_+ \quad (30)$$

a fixed conductance function.

Given (G, c) we introduce the Hilbert space $\mathcal{H}_E$ and the operator Δ as in (12), (13) and (14) above. Note that both depend on the choice of the function c in (30).

(a) Set

$$\text{Harm} := \{h \in \mathcal{H}_E \mid \Delta h = 0\}; \quad (31)$$

the finite-energy harmonic functions, “Harm” is short for “harmonic”. Note that functions h in Harm may be unbounded as functions on G^0 .

Setting $\text{Fin} :=$ the closed span of $\{\delta_x | x \in G^0\}$ in $\mathcal{H}_E$, it is easy to see that

$$\text{Harm} = \mathcal{H}_E \ominus \text{Fin}, \quad (32)$$

and

$$\text{Fin} = \mathcal{H}_E \ominus \text{Harm}, \quad (33)$$

i.e., that the decomposition

$$\mathcal{H}_E = \text{Fin} \oplus \text{Harm} \quad (34)$$

holds; see [22].

(b) Set

$$\text{Def} := \{u \in \mathcal{H}_E | \Delta u = -u\}; \quad (35)$$

the finite-energy deficiency vectors for the operator Δ .

(The reader will notice that solutions $u \neq 0$ to (35) appear to contradict the estimates (22) and (29); but this is not so. The deeper explanation lies in the theory of *deficiency-indices*, and *deficiency-subspaces* for unbounded linear operators in Hilbert space.) The notation “Def” is short for deficiency-subspace.

III. COMPUTATIONS

While the analysis of the two closed subspaces from Definition 2.5 (the finite energy harmonic functions, and the functions in the defect-space, or deficiency-space, for the Laplacian) will be carried out in general (sections VI and VII), it will be useful to work them out in a particular family of special cases; examples. At a first glance, these examples may indeed appear rather special, but we will show in section VII that they have wider use in our analysis of the “boundary at infinity.” The special cases discussed here will further show that the abstract spaces and operators from section II allow for explicit computations.

Example 3.1. *In an infinite weighted graph (G, c) take for vertex-set*

$$G^0 = \{0\} \cup \mathbb{Z}_+ = \{0, 1, 2, \dots\}, \quad (36)$$

edges, nearest neighbors,

$$\text{Nbh}_G(0) = \{1\}, \quad (37)$$

and

$$\text{Nbh}_G(x) = \{x-1, x+1\} \text{ if } x \in \mathbb{Z}_+. \quad (38)$$

Pick a function μ on $\mathbb{Z}_+$, and set

$$c(x-1, x) = : \mu(x) \text{ if } x \in \mathbb{Z}_+. \quad (39)$$

We will be interested in an *extended* version of the example when $G^0 = \mathbb{Z}$, and

$$\text{Nbh}_G(x) = \{x-1, x+1\} \quad (40)$$

for all $x \in \mathbb{Z}$. In this case, we will extend (39) by symmetry, as follows:

$$c(x-1, x) = : \mu(x) \text{ if } x \in \mathbb{Z}_+, \quad (41)$$

and

$$c(x, x+1) = \mu(|x|) \text{ if } x \in \mathbb{Z}_-. \quad (42)$$

To distinguish the two cases we will denote the first one (G_+, μ) , and the second (G, μ) .

It turns out that there are important distinctions between the two, relative to the two subspaces Harm and Def introduced above.

The matrix form of the operator Δ is as follows in the two cases:

$$\begin{pmatrix} \mu(1) & -\mu(1) & 0 & 0 & 0 & \cdots & \rightarrow \infty \\ -\mu(1) & \mu(1)+\mu(2) & -\mu(2) & 0 & 0 & & \\ 0 & -\mu(2) & \mu(2)+\mu(3) & -\mu(3) & 0 & & \\ 0 & 0 & -\mu(3) & \mu(3)+\mu(4) & -\mu(4) & & \\ 0 & 0 & 0 & -\mu(4) & \mu(4)+\mu(5) & \cdots & \\ \vdots & \vdots & \vdots & & & \cdots & \\ \downarrow & & & & & & \\ \infty & & & & & & \end{pmatrix}$$

TABLE 1. Matrix of Δ in (G_+, μ)

$$\begin{pmatrix} & 0 & & & & & & \\ & 0 & 0 & 0 & \cdots & \cdots & & \\ \ddots & -\mu(4) & 0 & 0 & 0 & \cdots & \cdots & \\ & \mu(3)+\mu(4) & -\mu(3) & 0 & 0 & 0 & \cdots & \\ & -\mu(3) & \mu(2)+\mu(3) & -\mu(2) & 0 & 0 & 0 & \cdots \\ & 0 & -\mu(2) & \mu(1)+\mu(2) & -\mu(1) & 0 & 0 & 0 \\ \cdots & 0 & 0 & -\mu(1) & 2\mu(1) & -\mu(1) & 0 & 0 \\ \cdots & 0 & 0 & 0 & -\mu(1) & \mu(1)+\mu(2) & -\mu(2) & 0 \\ & \cdots & \cdots & 0 & 0 & -\mu(2) & \mu(2)+\mu(3) & -\mu(3) \\ & & & \cdots & \cdots & 0 & -\mu(3) & \mu(3)+\mu(4) \\ & & & & & & & \ddots \\ & & & & & & 0 & 0 \end{pmatrix}$$

TABLE 2. Matrix of Δ in (G, μ)

Let $M \in \mathbb{R}_+$; suppose $M > 1$, and set $m = 1/M$. For the conductance function μ in (41) and (42) we set

$$\mu(x) := M^x = \exp(x \ln M), \quad x \in \mathbb{Z}_+. \quad (43)$$

Theorem 3.2. *In the examples with μ as in (43), we have the following*

	(G_+, μ)	(G, μ)
Harm	¹ zero	² one-dimensional
Def	³ one-dimensional	⁴ dimension zero or one; see details below!

TABLE 3. The four possibilities marked in the table will be proved in details below in parts 1 through 4.

Proof. It will be convenient for us to organize the proof into the four parts indicated in Table 3.

Part 1. With the definitions as given, we consider the possible solutions $h: \{0\} \cup \mathbb{Z}_+ \rightarrow \mathbb{C}$ to the equation

$$\Delta h = 0. \quad (44)$$

Hence

$$0 = \mu(1)(h(0) - h(1)), \quad (45)$$

and

$$0 = \mu(x)(h(x) - h(x-1)) + \mu(x+1)(h(x) - h(x+1)) \quad \text{for } x \in \mathbb{Z}_+. \quad (46)$$

Setting

$$(\delta h)(x) := h(x) - h(x-1), \quad (47)$$

equations (45) and (46) then take the following form:

$$\mu(1)(\delta h)(1) = 0, \quad (48)$$

and

$$\mu(x)(\delta h)(x) = \mu(x+1)(\delta h)(x+1) \quad \text{for all } x \in \mathbb{Z}_+. \quad (49)$$

Since $\mu > 0$, clearly the last two equations imply $\delta h \equiv 0$. So

$$h(x) = h(0) + \delta(1) + \cdots + (\delta h)(x) = h(0),$$

and h must be the constant function. But $\mathcal{H}_E$ is obtained by modding out with the constants, so $h = 0$ in $\mathcal{H}_E$.

Part 2. In this case, $G^0 = \mathbb{Z}$, and we set $h(0) = 0$, and

$$h(-x) = -h(x), \quad x \in \mathbb{Z}. \quad (50)$$

In that case,

$$\begin{aligned} 0 &= (\Delta h)(0) = \mu(1)(h(0) - h(-1) + h(0) - h(1)) \\ &= \mu(1)(2h(0) - h(-1) - h(1)), \end{aligned}$$

so conditions (50) work. This together with (49) then determine h on $\mathbb{Z}$.

Pick some constant $t \in \mathbb{R}_+$, and set

$$\mu(x) \delta h(x) = t \text{ for all } x \in \mathbb{Z}_+; \quad (51)$$

and then

$$\begin{aligned} h(x) &= \delta h(1) + \delta h(2) + \cdots + (\delta h)(x) \\ &= t(\mu(1)^{-1} + \mu(2)^{-1} + \cdots + \mu(x)^{-1}). \end{aligned}$$

With conductance $\mu(x) = M^x$, and resistance $\xi := M^{-1}$, we get

$$\begin{aligned} h(x) &= t(\xi + \xi^2 + \cdots + \xi^x) \\ &= t\xi \frac{1 - \xi^x}{1 - \xi} \xrightarrow{x \rightarrow \infty} \frac{t\xi}{1 - \xi}. \end{aligned} \quad (52)$$

Moreover,

$$\begin{aligned} \mathcal{E}(h) &= 2 \sum_{x=1}^{\infty} \mu(x) ((\delta h)(x))^2 \\ &= 2 \sum_{x=1}^{\infty} M^x (t\xi^x)^2 \\ &= 2t^2 \sum_{x=1}^{\infty} \xi^x = \frac{2t^2\xi}{1 - \xi} < \infty. \end{aligned}$$

□

Part 3. Now the “defect equation” is

$$\Delta u = -u, \quad u \in \mathcal{H}_E, \quad (53)$$

starting with

$$(\Delta u)(0) = \mu(1)(u(0) - u(1)) = -u(0);$$

so

$$(\delta u)(1) = u(1) - u(0) = \xi u(0),$$

or

$$u(1) = (1 + \xi)u(0). \quad (54)$$

Equation (53) for $x \in \mathbb{Z}_+$ yields

$$\mu(x+1)(\delta u)(x+1) = \mu(x)(\delta u)(x) + u(x). \quad (55)$$

Now set $u(0) = \lambda$. For $x = n \in \mathbb{Z}_+$, we now define two sequences of polynomials as follows

$$(\delta u)(n) := \xi^n p_n(\xi) \lambda, \quad (56)$$

$$u(n) := q_n(\xi) \lambda, \quad (57)$$

where

$$\xi = M^{-1}, \text{ and } \mu(n) = M^n. \quad (58)$$

It follows that u is a multiple of the function

$$u_1(x) := q_x(\xi), \quad x \in 0 \cup \mathbb{Z}_+. \quad (59)$$

Specifically, $q_0 = 1$, $p_1 = 1$, $q_1 = 1 + \xi$; and

$$p_{n+1} = p_n + q_n \quad (60)$$

$$q_{n+1} = q_n + \xi^{n+1} p_{n+1} = \xi^{n+1} p_n + (1 + \xi^{n+1}) q_n. \quad (61)$$

In matrix form:

$$\begin{pmatrix} p_n(\xi) \\ q_n(\xi) \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ \xi^n & 1 + \xi^n \end{pmatrix} \begin{pmatrix} 1 & 1 \\ \xi^{n-1} & 1 + \xi^{n-1} \end{pmatrix} \cdots \begin{pmatrix} 1 & 1 \\ \xi^2 & 1 + \xi^2 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ \xi & 1 + \xi \end{pmatrix} \begin{pmatrix} 0 \\ 1 \end{pmatrix}. \quad (62)$$

The polynomials may well be of independent interest, and we will need the first few in the infinite string:

n	$p_n(\xi)$	$q_n(\xi)$
$n = 1$	1	$1 + \xi$
$n = 2$	$2 + \xi$	$\xi^2 + (1 + \xi^2)(1 + \xi)$ $= 1 + \xi + 2\xi^2 + \xi^3$
$n = 3$	$3 + 2\xi + 2\xi^2 + \xi^3$	$\xi^3(2 + \xi) + (1 + \xi^3)(1 + \xi + 2\xi^2 + \xi^3)$ $= 1 + \xi + 2\xi^2 + 4\xi^3 + 2\xi^4 + 2\xi^5 + \xi^6$

TABLE 4.

Lemma 3.3. *The first and the last terms in the polynomials $p_n(\xi)$ and $q_n(\xi)$ are as indicated in the following formulas:*

$$p_n(\xi) = n + (n-1)\xi + \cdots + \xi^{\frac{(n-1)n}{2}}, \quad (63)$$

and

$$q_n(\xi) = 1 + \xi + \cdots + \xi^{\frac{n(n+1)}{2}}. \quad (64)$$

So the degree of $q_n(\xi)$ is $\frac{n(n+1)}{2}$.

Proof. Note that Table 4 already suggests the start of an induction proof. Now suppose (63) and (64) hold up to n , i.e., $p_n(0) = n$; and that both $p_n(\xi)$ and $q_n(\xi)$ have leading coefficients one.

With the use of (60) and (61) we then get

$$\begin{aligned} p_{n+1}(0) &= p_n(0) + q_n(0) \\ &= n + 1, \end{aligned}$$

which is the next step in the induction. Now apply the same argument to

$$\left. \frac{d}{d\xi} p_{n+1} \right|_{\xi=0},$$

and the result in (63) for the next coefficient follows.

Setting $\xi = 0$ in

$$q_{n+1}(\xi) = \xi^{n+1} p_n(\xi) + (1 + \xi^{n+1}) q_n(\xi) \quad (65)$$

and using the induction hypothesis, we get $q_{n+1}(0) = q_n(0) = 1$.

We now turn to the leading coefficient in $q_{n+1}(\xi)$. As before, we use the induction hypothesis, and (65). We get

$$\begin{aligned} q_{n+1}(\xi) &= \xi^{n+1} \left(n + \dots + \xi^{\frac{(n-1)n}{2}} \right) + (1 + \xi^{n+1}) \left(1 + \dots + \xi^{\frac{n(n+1)}{2}} \right) \\ &= 1 + \dots + \xi^{\frac{(n+1)(n+2)}{2}}. \end{aligned}$$

This completes the induction proof.

In other words, the degree of $q_n(\xi)$ is

$$1 + 2 + 3 + \dots + n = \frac{n(n+1)}{2};$$

and each $q_n(\xi)$ has leading coefficient one. $\square$

Proposition 3.4.

(a) *The two generating functions*

$$P(X, \xi) := \sum_{n=1}^{\infty} p_n(\xi) X^n, \quad (66)$$

and

$$Q(X, \xi) := \sum_{n=0}^{\infty} q_n(\xi) X^n \quad (67)$$

satisfy

$$X(P(X, \xi) + Q(X, \xi)) = P(X, \xi) \quad (68)$$

and

$$Q(X, \xi) = 1 + XQ(X, \xi) + P(\xi X, \xi). \quad (69)$$

(b) Here we have picked $\xi \in (0, 1)$; and we note that, in the first variable, $Q(X, \xi)$ has radius of convergence 1, while $P(X, \xi)$ has radius of convergence $\sqrt{\xi}$.

Proof. Multiplying through by X^n in

$$p_{n+1}(\xi) = p_n(\xi) + q_n(\xi),$$

and using

$$\begin{pmatrix} p_0 \\ q_0 \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \quad (70)$$

we arrive at the first formula (68) in the statement of the proposition.

For the proof of (69) we again multiply through by X^{n+1} , now in

$$q_{n+1}(\xi) = q_n(\xi) + \xi^{n+1} p_{n+1}(\xi).$$

After adding up the terms with “ $\sum_{n=0}^{\infty} \dots$ ” and using (70), we arrive at the desired conclusion (69).

We now turn to the radii of convergence:

Since we already established that

$$\sum_{n=1}^{\infty} \xi^n p_n(\xi)^2 < \infty,$$

it follows that there is a finite constant C such that

$$p_n(\xi) \leq C \xi^{-\frac{n}{2}}; \quad (71)$$

and we conclude that $X \mapsto P(X, \xi)$ has radius of convergence $\sqrt{\xi}$ as claimed.

But further note that $n \mapsto q_n(\xi)$ is bounded, as a consequence of the estimate (71). Hence we conclude that $X \mapsto Q(X, \xi)$ has radius of convergence = 1. $\square$

The purpose of the previous discussion is to find the deficiency vector u , i.e., the solution u in $\mathcal{H}_E$ in a random walk model with $p(n, n+1) = \frac{M}{1+M}$, $p(n, n-1) = \frac{1}{1+M}$ where $M > 1$. We now turn to the corresponding generating functions

$$Q\left(X, \frac{1}{M}\right) = \sum_{n=1}^{\infty} u(n) X^n.$$

Corollary 3.5. *The generating function $X \mapsto P(X, \xi)$ from (66) in Proposition 3.4 has the representation*

$$P(X, \xi) = \sum_{n=1}^{\infty} \frac{\xi^{\frac{n(n+1)}{2}} X^n}{(1-X)^2 (1-\xi X)^2 (1-\xi^2 X)^2 \cdots (1-\xi^n X)^2}. \quad (72)$$

Proof. This is an application of (66) and (67): Eliminate $Q(X, \xi)$ and iterate the substitution. $\square$

Corollary 3.6. *The generating function for u itself is as follows:*

$$\begin{aligned} Q(X, \xi) &= \sum_{n=0}^{\infty} u(n, \xi) X^n \\ &= \frac{1}{1-X} \left(1 + \sum_{n=1}^{\infty} \frac{\xi^{\frac{n(n+1)}{2}} X^n}{(1-\xi X)^2 \cdots (1-\xi^n X)^2} \right), \end{aligned}$$

where

$$\prod_{k=1}^{\infty} (1 - \xi^k X) = \exp \left(-\frac{\xi X}{1-X} \right).$$

Proof. Combine the previous formulas. $\square$

In the next section, we take up a number of dynamics related issues concerning these polynomials. Below we are concerned with the proof of the following:

Lemma 3.7. *For every $\xi \in (0, 1)$, there is an $m \in \mathbb{N}$ such that*

$$p_x(\xi) \leq x^m, \quad (73)$$

and

$$q_x(\xi) \leq (x+1)^m - x^m, \text{ for all } x \in \mathbb{Z}_+, \quad (74)$$

where $m = m(\xi)$ depends on ξ .

Proof. Table 4 makes clear the start of an induction of (73) and (74). Now suppose m has been chosen such that (73) and (74) hold up to x .

Then

$$\begin{aligned} p_{x+1}(\xi) &= p_x(\xi) + q_x(\xi) \\ &\leq x^m + (x+1)^m - x^m \\ &= (x+1)^m. \end{aligned}$$

As a result, we get

$$\begin{aligned} q_{x+1}(\xi) &= q_x(\xi) + \xi^{x+1} p_{x+1}(\xi) \\ &\leq (x+1)^m - x^m + \xi^{x+1} (x+1)^m. \end{aligned} \quad (75)$$

Now in the next step, we adjust m such that

$$(x+1)^m - x^m + \xi^{x+1} (x+1)^m \leq (x+2)^m - (x+1)^m.$$

We rewrite this:

$$\begin{aligned} 2 + \xi^{x+1} &\leq \left(1 - \frac{1}{x+1}\right)^m + \left(1 + \frac{1}{x+1}\right)^m \\ &\leq 2 \cdot \left(1 + \binom{m}{2} \left(\frac{1}{x+1}\right)^2\right); \end{aligned}$$

or

$$(x+1)^2 \xi^{x+1} \leq m(m-1). \quad (76)$$

But

$$\max_{t \in \mathbb{R}_+} t^2 \xi^t = \left(\frac{2}{\ln \xi}\right)^2 e^{-2}. \quad (77)$$

It follows that (76) holds if

$$|\ln \xi| > \frac{2}{e\sqrt{m(m-1)}},$$

and so

$$M > \exp\left(\frac{2}{e\sqrt{m(m-1)}}\right). \quad (78)$$

Since the limit on the RHS in (78) is 1 as $m \rightarrow \infty$, and $M > 1$ is fixed, it follows that m can be adjusted to $\xi = 1/M$ such that (78) holds. With this choice in fact (74) will be satisfied for all $x \in \mathbb{Z}_+$. To see this, apply (77) to $t = x+1$. $\square$

Lemma 3.8. *The solution u from Part 3 in Table 3 satisfies*

$$\mathcal{E}(u) < \infty \quad (79)$$

Proof.

$$\begin{aligned} \mathcal{E}(u) &= \sum_{x \in \mathbb{Z}_+} \mu(x) ((\delta u)(x))^2 \\ &=_{(\text{by Lemma 3.8})} \sum_{x \in \mathbb{Z}_+} M^x (\xi^x p_x(\xi))^2 \\ &\leq \sum_{x \in \mathbb{Z}_+} x^{2m} \xi^x < \infty. \end{aligned}$$

$\square$

This finishes the proof of Part Three in Theorem 3.2.

Part 4. Here the model is $(\mathbb{Z}, \mu)$ when the function μ satisfies (41) and (42). As a result the solution u to $\Delta u = -u$ considered in the theorem satisfies

$$u(x) = u(-x), \quad x \in \mathbb{Z}. \quad (80)$$

We get

$$(\Delta u)(0) = M(2u(0) - u(1)) = -2M(\delta u)(1) = -u(0).$$

Hence

$$(\delta u)(1) = \left(\frac{\xi}{2}\right) u(0), \quad (81)$$

and

$$u(1) = \left(1 + \frac{\xi}{2}\right) u(0).$$

Now $u(x)$ is the second component in the vector

$$\begin{aligned} \begin{pmatrix} * \\ u(x) \end{pmatrix} &= \begin{pmatrix} 1 & 1 \\ \xi^x & 1 + \xi^x \end{pmatrix} \begin{pmatrix} 1 & 1 \\ \xi^{x-1} & 1 + \xi^{x-1} \end{pmatrix} \\ &\quad \cdots \begin{pmatrix} 1 & 1 \\ \xi^2 & 1 + \xi^2 \end{pmatrix} \begin{pmatrix} \frac{1}{2} \\ 1 + \frac{\xi}{2} \end{pmatrix} u(0). \end{aligned}$$

□

IV. THE POLYNOMIALS $p_n(z)$ AND $q_n(z)$, $n = 1, 2, \dots$

We begin with some technical lemmas, and we further observe that the examples from section III indeed have a more general flavor: for example (Proposition 5.1), they may be derived from a standard random walk model. We will do the computations here just for a one-dimensional walk, but the basic idea carries over much more generally as we show in the next section.

OPTIMAL ESTIMATES. In the previous section, we considered $M > 1$, $\xi = 1/M$, and two sequences $p_n(\xi)$, $q_n(\xi)$ for $n = 1, 2, 3, \dots$. In the proof of Theorem 3.2, we showed the following:

Proposition 4.1. *If*

$$M > \exp\left(e^{-1}\sqrt{2/3}\right), \quad (82)$$

then

$$p_n(\xi) \leq n^3 \text{ for all } n \in \mathbb{Z}_+. \quad (83)$$

In the following sense, this is not optimal; in fact, there is a finite constant B such that we get linear bounds in both directions. Specifically:

$$n \leq p_n(\xi) \leq 1 + B \left(\frac{\sqrt{\xi}}{1 - \sqrt{\xi}} \right) n. \quad (84)$$

Proof. We will establish the existence of $B \in \mathbb{R}_+$ by induction; and the size of B will follow from the *a priori* estimates to follow. The induction begins with an inspection of Table 4 in section III.

Lemma 4.2. *For every $\xi \in (0, 1)$, the following finite limit exists:*

$$Q(\infty, \xi) := \lim_{n \rightarrow \infty} q_n(\xi); \quad (85)$$

the limit is monotone, and $Q(\infty, \xi) > 1$.

Proof. Let $u = u_\xi$ be the function from Lemma 3.7.

We established that

$$\mathcal{E}(u_\xi) = \sum_{n=1}^{\infty} \xi^n p_n(\xi)^2 < \infty$$

where $(\delta u_\xi)(n) = \xi^n p_n(\xi)$. If $\xi^n p_n(\xi)^2 \leq A$, $\forall n \in \mathbb{Z}_+$, for some fixed constant A ; then

$$(\delta u_\xi)(n) \leq \sqrt{A} \xi^{n/2}. \quad (86)$$

Since

$$u_\xi(n) = 1 + (\delta u_\xi)(1) + \cdots + (\delta u_\xi)(n), \quad (87)$$

we get

$$\begin{aligned} u_\xi(n) &\leq 1 + \sqrt{A\xi} \frac{1 - \xi^{n/2}}{1 - \sqrt{\xi}} \\ &< 1 + \frac{\sqrt{A\xi}}{1 - \sqrt{\xi}}. \end{aligned}$$

Since

$$u_\xi(1) < u_\xi(2) < \cdots < u_\xi(n) < u_\xi(n+1) < \cdots, \quad (88)$$

the desired conclusion follows, and

$$Q(\infty, \xi) \leq 1 + \frac{\sqrt{A\xi}}{1 - \sqrt{\xi}}. \quad (89)$$

We will see below that

$$p_n(\xi) = 1 + \sum_{x=1}^{n-1} q_x(\xi),$$

so

$$p_n(\xi) \leq 1 + \left(\frac{\sqrt{A\xi}}{1 - \sqrt{\xi}} \right) n.$$

The estimate in the RHS in (87) now follows.

Returning to $p_n(\xi)$, we have

$$\begin{aligned} p_n(\xi) &= p_{n-1}(\xi) + q_{n-1}(\xi) \\ &= q_{n-1}(\xi) + q_{n-2}(\xi) + \cdots + q_1(\xi) + p_1(\xi) \end{aligned} \quad (90)$$

where

$$\begin{pmatrix} p_1(\xi) \\ q_1(\xi) \end{pmatrix} = \begin{pmatrix} 1 \\ 1 + \xi \end{pmatrix}.$$

Since

$$q_x(\xi) \geq 1 \text{ for } x \in \mathbb{Z}_+,$$

(90) implies

$$p_n(\xi) \geq n,$$

which is the remaining lower boundary on the LHS in (87). $\square$

$\square$

V. A RANDOM WALK MODEL

In this section we resume our analysis of the general case: How does the operator theory throw light on “the boundary at infinity” as it was made precise in section III? Part of the answer lies in a transfer operator studied in Lemma 5.6 below.

For the weighted graphs (G, c) in section II with $c : G^1 \rightarrow \mathbb{R}_+$ representing conductance we may introduce

$$c(x) = \sum_{y \sim x} c(x, y)$$

and

$$p(x, y) := \frac{c(x, y)}{c(x)}. \quad (91)$$

Then

$$\sum_{y \sim x} p(x, y) = 1 \quad (92)$$

and the function $p(x, y)$ represents a system of transition probabilities. They in turn determine a random walk on the set G^0 of all vertices. If ω is a path of a “walker”, and $X_n(\omega)$ is the location in G^0 at time n , then

$$\text{Prob}(\{\omega | X_n(\omega) = x, X_{n+1}(\omega) = y\}) = p(x, y) \text{ for all } (x, y) \in G^1, \quad (93)$$

where “Prob” is short for “probability.” This is a *reversible* random walk in that

$$c(x)p(x, y) = c(y)p(y, x) \text{ for all } x \sim y. \quad (94)$$

Proposition 5.1. *In the model $(\mathbb{Z}_+, M^x)$ in Theorem 3.2, the transition probability from 0 to 1 is 1, and from n to $n + 1$, it is the fixed quotient $M/1+M$ for all $n \in \mathbb{Z}_+$. Specifically, a move to the right:*

$$p(n, n + 1) = \frac{M}{1 + M};$$

and

$$p(n, n-1) = \frac{1}{1+M}$$

equals the probability of a move to the left when starting at the vertex n .

Proof. Since the edges in $\mathbb{Z}_+ \cup \{0\}$ are just the nearest neighbors, we get

$$\begin{aligned} p(0, 1) &= 1, \text{ and} \\ p(n, n+1) &= \frac{M^{n+1}}{M^n + M^{n+1}} \\ &= \frac{M^{n+1}}{M^n \cdot (1 + M)} \\ &= \frac{M}{1 + M}. \end{aligned}$$

□

Proposition 5.2. *In the model on $\mathbb{Z}$ with the following*

$$c(0, 1) = A, c(n-1, n) = A^n, \quad n \in \mathbb{Z}_+;$$

and

$$c(-1, 0) = B$$

and

$$c(-n, -n+1) = B^n$$

we get the following transition probabilities:

$$\begin{aligned} p(0, 1) &= \frac{A}{A+B}, \\ p(0, -1) &= \frac{B}{A+B}, \\ p(n, n+1) &= \frac{A}{1+A}, \\ p(n, n-1) &= \frac{1}{1+A}, \text{ for } n \in \mathbb{Z}_+; \text{ and} \\ p(-n, -n-1) &= \frac{B}{1+B}, \text{ and} \\ p(-n, -n+1) &= \frac{1}{1+B}. \end{aligned}$$

Proof. Left to the reader.

□

The significance of the deficiency vectors u studied in section II is that the presence of a non-zero deficit eigenspace

$$\text{Def} := \{u \in \mathcal{H}_E \mid \Delta u = -u\} \quad (95)$$

offers a quantitative measure of states at “infinity” in the model.

To make this precise, we return to the operator theory for the operator Δ with its dense domain $\mathcal{D}$ in $\mathcal{H}_E$; see Proposition 2.4 for details.

Definition 5.3. *The domain of the adjoint operator Δ^* is defined as follows*

$$\text{dom}(\Delta^*) = \{u \in \mathcal{H}_E \mid \text{s.t. } \exists C > \infty \text{ with } |\langle u, \Delta\varphi \rangle| \leq C \|\varphi\|, \forall \varphi \in \mathcal{D}\} \quad (96)$$

If $u \in \text{dom}(\Delta^)$, set $\Delta^*u = v$ if*

$$\langle v, \varphi \rangle = \langle u, \Delta\varphi \rangle \text{ holds for all } \varphi \in \mathcal{D}. \quad (97)$$

We say that Δ is essentially selfadjoint if its closure Δ^{clo} is selfadjoint, i.e., if $\Delta^{clo} = \Delta^*$.

Now Δ is semibounded on its domain $\mathcal{D}$ in $\mathcal{H}_E$ by Proposition 2.4; and it thus follows from a theorem by von Neumann [11] that Δ is essentially selfadjoint iff the corresponding deficiency space Def is zero. To make the connection between the operator theory and the solutions u studied in section III, we will need the following.

Lemma 5.4. *Let $\mathcal{H}_E, \Delta, \mathcal{D}$ and Δ^* be as in Definition 5.3. If $u \in \text{dom}(\Delta^*)$, then*

$$(\Delta^*u)(x) = \sum_{y \sim x} c(x, y) (u(x) - u(y)). \quad (98)$$

Proof. Note that the assumptions in the lemma include the following two assertions:

- (i) $u \in \mathcal{H}_E$, and
- (ii) $\Delta^*u \in \mathcal{H}_E$.

Now let $y \in G^0 \setminus (0)$, and compute

$$\begin{aligned} \langle v_y, \Delta^*u \rangle_E &= \langle \Delta v_y, u \rangle_E \\ &= \langle \delta_y - \delta_o, u \rangle_E \\ &= \sum_{z \sim y} c(y, z) (u(y) - u(z)) - \sum_{z \sim o} c(o, z) (u(o) - u(z)) \\ &= \left\langle v_y(\cdot), \sum_z c(\cdot, z) (u(\cdot) - u(z)) \right\rangle_E. \end{aligned}$$

Since the family (v_y) spans the dense subspace $\mathcal{D}$ in $\mathcal{H}_E$, the desired formula (98) now follows. Note that (98) is the identity of two vectors in the energy Hilbert space $\mathcal{H}_E$. $\square$

For the two-sided random walk on $\mathbb{Z}$ with conductances given in Figure 1 below, we are now ready to compute the deficit space Def in $\mathcal{H}_E$.




Figure 1. Conductance.

Proposition 5.5. *The deficit vectors $u \in \mathcal{H}_E$ solving $\Delta^*u = -u$ in the $A-B$ model on $\mathbb{Z}$ given in Figure 1 are determined by the following system:*

$$(A + B + 1)u(0) = Au(1) + Bu(-1) \quad (99)$$

and

$$p_1(\alpha) + p_1(\beta) = 1, \quad (100)$$

where $\alpha = 1/A$, and $\beta = 1/B$.

Setting $\mu(0) = 1$, we then obtain the following recursive determination: If $n \in \mathbb{Z}_+$, then

$$\mu(n) = q_n(\alpha), u(-n) = q_n(\beta)$$

with

$$\begin{pmatrix} p_n(\xi) \\ q_n(\xi) \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ \xi^n & 1 + \xi^n \end{pmatrix} \cdots \begin{pmatrix} 1 & 1 \\ \xi^2 & 1 + \xi^2 \end{pmatrix} \begin{pmatrix} 1 & 1 \\ \xi & 1 + \xi \end{pmatrix} \begin{pmatrix} 0 \\ 1 \end{pmatrix} \quad (101)$$

for $\xi = \alpha$ on the $\mathbb{Z}_+$ side, and $\xi = \beta$ on the $\mathbb{Z}_-$ side.

Proof. By the lemma, we get the following identities:

$$A(\delta u)(1) + B(\delta u)(-1) = u(0).$$

With $u(0) = 1$, this amounts to equations (99) and (100).

Using

$$u(n) = u(n-1) + (\delta u)(n), \quad n \in \mathbb{Z}_+,$$

we get

$$u(n) = q_n(\alpha)$$

with

$$q_n(\alpha) = q_{n-1}(\alpha) + \alpha^n p_n(\alpha).$$

With

$$u(-n) = u(-n+1) + (\delta u)(-n),$$

we get ($n \in \mathbb{Z}_+$):

$$u(-n) = q_n(\beta)$$

with

$$q_n(\beta) = q_{n-1}(\beta) + \beta^n p_n(\beta).$$

The resulting formulas (101) (in matrix form) now follow from this; and as a result the solution u is determined on all of $\mathbb{Z}$, with the normalizations $u(0) = 1$, and (100). $\square$

Lemma 5.6. *Let (G, c) be a weighted graph with probabilities*

$$p(x, y) := \frac{c(x, y)}{c(x)}, \quad (102)$$

and set

$$(Tf)(x) = \sum_y p(x, y) f(y); \quad (103)$$

T for transfer operator.

(a) *Then*

$$Tf = f - \frac{1}{c} \Delta f. \quad (104)$$

A vector $f \in \mathcal{H}_E$ is in Def if and only if

$$Tf = \left(1 + \frac{1}{c}\right) f \quad (105)$$

where

$$c(x) = \sum_{y \sim x} c(x, y).$$

(b) *In the example of Proposition 5.1 ($G^0 = \{0\} \cup \mathbb{Z}_+$),*

$$(Tf)(x) = \frac{1}{1+M} f(x-1) + \frac{M}{1+M} f(x+1), \text{ for all } x \in G^0. \quad (106)$$

(c) *In general if $f \in \text{Def}$ is bounded, then the pointwise limit*

$$\lim_{k \rightarrow \infty} (T^k f)(x) =: h(x) \quad (107)$$

exists, and h is harmonic.

Proof. (a) We have

$$\begin{aligned} (\Delta f)(x) &= \sum_y c(x, y) (f(x) - f(y)) \\ &= c(x) f(x) - \sum_y c(x, y) f(y) \\ &= c(x) (f(x) - (Tf)(x)), \end{aligned}$$

and (104) follows.

(b) follows from (a) and substitution into (102), i.e.,

$$c(n-1, n) = M^n, \text{ for } n \in \mathbb{Z}_+.$$

(c) follows from (a) and (b).

In particular if the Δ a constant $A \in \mathbb{R}_+$ such that $f(x) \leq A$ for all $x \in G^0$, then apply T to the positive function $A - f$, and we get $A \geq T^k f$ pointwise on G^0 . Since $Tf = (1 + \frac{1}{c}) f$ we get

$$f \leq Tf \leq T^2 f \leq \dots \leq T^k f \leq \dots \leq A.$$

Therefore the limit $h(x)$ in (107) exists if and only if $f \in \text{Def}$ is a bounded function.

The following is then immediate:

$$\Delta h = 0 \Leftrightarrow Th = h.$$

□

VI. THE SPACE Def AND BOUNDARY CONDITIONS

In this section, we discuss the distinction between the operator and Δ considered in the Hilbert space $\mathcal{H}_E$ as compared with $\ell^2 := \ell^2(G^0)$. As before (G, c) is a given weighted graph with $c : G^1 \rightarrow \mathbb{R}_+$ representing conductances on the edges; and G^0 density the set of vertices. The corresponding Laplace operator is discussed in sections II and III, and we recall that it depends on the choice of function c .

When Δ is viewed as an operator in $\mathcal{H}_E$, we take as its domain

$$\mathcal{D}_V := \text{span} \{v_x | x \in G^0 \setminus \{o\}\} \quad (108)$$

where o is fixed in G^0 .

When Δ is viewed as an operator in ℓ^2 , we take as its domain

$$\mathcal{D}_E := \text{all finitely supported functions on } G^0. \quad (109)$$

To stress the distinction between the two cases $\mathcal{H}_E$ and ℓ^2 , we shall use the following terminology, $(\Delta, \mathcal{D}_V, \mathcal{H}_E)$ and $(\Delta, \mathcal{D}_F, \ell^2)$. The corresponding inner products will be distinguished with subscripts $\langle \cdot, \cdot \rangle_E$

for $\mathcal{H}_E$, and $\langle \cdot, \cdot \rangle_2$ or $\langle \cdot, \cdot \rangle_{\ell^2}$ for the second, i.e.,

$$\langle u, v \rangle_{\ell^2} := \sum_{x \in G^0} \overline{u(x)} v(x). \quad (110)$$

We note that *symmetry* and *semiboundedness* is satisfied for Δ with respect to the two inner products, i.e.,

$$\langle u, \Delta v \rangle_E = \langle \Delta u, v \rangle_E, \text{ and } \langle u, \Delta u \rangle_E \geq 0 \text{ for all } u, v \in \mathcal{D}_V; \quad (111)$$

as well as

$$\langle u, \Delta v \rangle_2 = \langle \Delta u, v \rangle_2, \text{ and } \langle u, \Delta u \rangle_2 \geq 0, \text{ for all } u, v \in \mathcal{D}_F. \quad (112)$$

Lemma 6.1. [22] *The operator $(\Delta, \mathcal{D}_F, \ell^2)$ is essentially selfadjoint; and so in particular, there are no non-zero solutions to the deficit equations in ℓ^2 : If $\Delta u = -u$, and $u \in \ell^2$, then $u = 0$.*

By contrast, we saw in sections III and IV, that

$$\Delta u = -u \quad (113)$$

has non-zero solutions in $\mathcal{H}_E$ for a rich class of unbounded functions c on G^1 . And so in particular, $(\Delta, \mathcal{D}_V, \mathcal{H}_E)$ is *not* essentially selfadjoint.

Proposition 6.2. *If $u \in \text{Def} \setminus \{0\}$, then the infinite sum*

$$S_2(u) := \sum_{x \in G^0} \overline{u(x)} (\Delta u)(x) \quad (114)$$

is divergent.

Proof. Recall that $\Delta u = \Delta^* u = -u$ if u is in Def and, as a result,

$$S_2(u) = - \sum_x |u(x)|^2.$$

But, by Lemma 6.1, the solution u to (113) cannot be in ℓ^2 and, as a result,

$$\sum_x |u(x)|^2 = \infty.$$

□

Theorem 6.3. *If $u \in \mathcal{H}_E \ominus \text{Def}$, then the sum $S_2(u)$ in (114) is finite and it has the following representation:*

There is a unique $v \in \text{dom}(\Delta_V^{\text{clo}})$ such that

$$u = v + \Delta_V^{\text{clo}} v,$$

and

$$\mathcal{E}(u) = S_2(u) + \mathcal{E}(P_{\text{Harm}} v), \quad (115)$$

where $\mathcal{E}$ denotes the energy-form, i.e., $\mathcal{E}(\cdot) := \|\cdot\|_{\mathcal{H}_E}$ and where P_{Harm} denotes the projection onto the harmonic functions h in $\mathcal{H}_E$, i.e.,

$$\text{Harm} := \{h \in \mathcal{H}_E \mid \Delta h = 0\}. \quad (116)$$

Proof. We begin with the closed subspace Harm in (116).

Lemma 6.4. ([22])

$$\begin{aligned} \text{Harm} &= \{h \in \mathcal{H}_E \mid h \perp \delta_x, \forall x \in G^0\} \\ &= \mathcal{H}_E \ominus \{\delta_x\}_{x \in G^0}; \end{aligned} \quad (117)$$

or setting

$$\text{Fin} := \text{the closed linear span of } (\delta_x)_{x \in G^0}. \quad (118)$$

Proof. We have for all $u \in \mathcal{H}_E$,

$$(\Delta u)(x) = \langle \delta_x, u \rangle_E, \quad (119)$$

and it follows that $\Delta u \equiv 0$ iff u satisfies the conditions in (117).

If P_{Harm} and P_{Fin} denote the respective projections onto the closed subspaces in (117) and (118), then

$$P_{\text{Harm}} + P_{\text{Fin}} = I \quad (120)$$

where I denotes the identity operator in $\mathcal{H}_E$. $\square$

Lemma 6.5. Let $v, w \in \mathcal{D}_v$; then

$$\sum_{x \in G^0} \overline{v(x)} (\Delta w)(x) = \langle P_{\text{Fin}} v, w \rangle_{\mathcal{H}_E}. \quad (121)$$

Proof. Introduce the sum notation $\sum \bar{v} \Delta w$ for the left hand side expression in (121). Then

$$\begin{aligned} \sum \overline{v(x)} (\Delta w)(x) &=_{(\text{by (119)})} \sum_x \overline{v(x)} \langle \delta_x, w \rangle_E \\ &= \left\langle \sum_x v(x) \delta_x, w \right\rangle_E \\ &=_{(\text{by (118)})} \langle P_{\text{Fin}} v, w \rangle_E. \end{aligned}$$

$\square$

Lemma 6.6. If $v \in \text{dom}(\Delta_V^{\text{clo}})$, then

$$\|v + \Delta v\|_E \geq \|v\|_E, \quad (122)$$

and so

$$u \longmapsto (I + \Delta)^{-1} u = v \quad (123)$$

is contractive on the closed subspace $\mathcal{H}_E \ominus \text{Def}$ in $\mathcal{H}_E$.

Proof. We will be using the operator theoretic properties (111) and (112) stated before Lemma 6.1. In particular, we use that if an operator Δ is hermitian symmetric and semibounded on a dense domain in Hilbert space, then so is its closure $\Delta^{clo} = \Delta^{**}$. We will be using the same symbol Δ also when referring to the closed operator.

Now for $v \in \text{dom}(\Delta^{clo})$, we have

$$\begin{aligned} \|v + \Delta v\|_E^2 &= \langle v + \Delta v, v + \Delta v \rangle_E \\ &= \|v\|_E^2 + \langle \Delta v, v \rangle_E + \langle v, \Delta v \rangle_E + \|\Delta v\|_E^2 \\ &\geq \|v\|_E^2 \text{ by (111).} \end{aligned}$$

□

Proof of Theorem 6.3 resumed. We have

$$\begin{aligned} \text{Def} &= N_{\mathcal{H}_E}(I + \Delta_v^*) \\ &= (R(I + \Delta^{clo}))^\perp. \end{aligned} \tag{124}$$

By virtue of Lemma 6.6, we further note that $R(I + \Delta^{clo})$ is closed in $\mathcal{H}_E$, and as a result

$$R(I + \Delta^{clo}) = \mathcal{H}_E \ominus \text{Def}. \tag{125}$$

With $u = v + \Delta v$, we now compute as follows. For the expression S_2 in equation (114) we have:

$$\begin{aligned} S_2(u) &= \sum_x \bar{u} \Delta u \\ &= \sum_x \bar{v} \Delta v + \sum_x \bar{v} \Delta^2 v + \sum_x \overline{\Delta v} \Delta v + \sum_x \overline{\Delta v} \Delta^2 v \\ &= (\text{by Lemma (6.5)}) \langle P_{\text{Fin}} v, v \rangle_E + \langle P_{\text{Fin}} v, \Delta v \rangle_E + \langle P_{\text{Fin}} \Delta v, v \rangle_E + \|\Delta v\|_E^2 \\ &= \|P_{\text{Fin}} v\|_E^2 + \langle v, \Delta v \rangle_E + \langle \Delta v, v \rangle_E + \|\Delta v\|_E^2 \\ &= \|P_{\text{Fin}} v\|_E^2 - \|v\|_E^2 + \|v + \Delta v\|_E^2 \\ &= (\text{by (119)}) - \|P_{\text{Harm}} v\|_E^2 + \|u\|_E^2 \\ &= -\mathcal{E}(P_{\text{Harm}} v) + \mathcal{E}(u). \end{aligned}$$

The desired conclusion (115) in the statement of the theorem now follows. □

VII. EMBEDDINGS OF GRAPHS AND OF HILBERT SPACES

In this section we show that mappings between graphs (for example inclusions of sub-graphs into an ambient super-graph) induce embeddings of the corresponding Hilbert spaces. This is functorial, and it

further allows localization. By this we mean embeddings of certain local “portions” of an ambient infinite graph G will produce intertwining operators in such a way that local spectral information is embedded into global. One advantage of this is that computations can more easily be carried out in the “local portions” of G .

Following the definitions in section II, we shall now consider pairs of weighted graphs (G, c_G) and (H, c_H) . For the first one, we have a vertex set G^0 and corresponding edges G^1 ; and similarly with H^0 and H^1 for the second graph. The conductance functions c_G and c_H are as follows:

$$c_G : G^1 \rightarrow \mathbb{R}_+, \text{ and } c_H : H^1 \rightarrow \mathbb{R}_+, \quad (126)$$

each satisfying the axioms from section II. The corresponding energy Hilbert spaces will be denoted $\mathcal{H}_E(G, c_G)$ and $\mathcal{H}_E(H, c_H)$; or simply $\mathcal{H}_E(G)$ and $\mathcal{H}_E(H)$ when the choice of conductance functions are clear from the context.

Definition 7.1. *Consider two functions*

$$G^0 \xrightarrow{\varphi} H^0, \quad (127)$$

and

$$G^0 \xrightarrow{\psi} \mathbb{R}_+. \quad (128)$$

We say that a pair of functions φ, ψ as in (127) and (128) is compatible if the operator

$$Tu := u \circ \varphi \quad (129)$$

is a well-defined isometry mapping $\mathcal{H}_E(H)$ into $\mathcal{H}_E(G)$; and if

$$(T\Delta_H u)(x) = \psi(x)(\Delta_G Tu)(x). \quad (130)$$

Definition 7.2. *(with examples!) Let (H, c_H) be the graph with vertex set $H^0 := \{0\} \cup \mathbb{Z}$, and nearest neighbor edges; we set*

$$c_H(n) := c_H(n-1, n) = 2^n, \text{ for all } n \in \mathbb{Z}_+. \quad (131)$$

Let (G, c_G) be the graph with vertex set $G^0 =$ the dyadic tree (see Figure 2) extending to $+\infty$ on the right.




Figure 2. The dyadic tree.

Points in G^0 are finite words in the finite alphabet (of bits) $\{0, 1\}$. We include the empty word, denoted $\emptyset$, on the extreme left in Figure 2.

For the edges G^1 , we take the nearest neighbors in G^0 . Hence if $x \in G^0 \setminus \{\emptyset\}$, then x has three nearest neighbors as follows: If $x = (x_1, x_2, \dots, x_n)$, $x_i \in \{0, 1\}$, then

$$\text{Nb}_G(x) := \{x^T, (x0), (x1)\} \quad (132)$$

where $x^T = (x_1, x_2, \dots, x_{n-1})$, $(x0) = (x_1, \dots, x_n, 0)$, and $(x1) = (x_1, \dots, x_n, 1)$; see also Figure 3.




Figure 3. Three nearest Neighbors.

The remaining case is

$$\text{Nbh}_G(\emptyset) = \{0, 1\}. \quad (133)$$

The reader may check by computation that the following two functions $\varphi : G^0 \rightarrow H^0$, and $\psi : G^0 \rightarrow \mathbb{R}_+$ constitute a compatible pair; i.e., that if T is defined as in (129) then it is *isometric* and ψ -intertwining, see (130). We set

$$\begin{aligned} \varphi(x_1 x_2 \cdots x_n) &= n \text{ if } x \neq \emptyset, \\ \varphi(\emptyset) &= 0, \text{ and} \\ \psi(x) &:= 2^{\varphi(x)}, \quad x \in G^0. \end{aligned} \quad (134)$$

We now turn to some applications dealing with harmonic functions of finite energy, and monopoles.

Definition 7.3. *Let (G, c) be a weighted graph with conductance function c ; and let Δ and $\mathcal{H}_E$ be the corresponding Laplace operator, and energy Hilbert space, respectively. Pick $o \in G^0$; and set*

$$\text{Mono}(G) := \{w \in \mathcal{H}_E \mid \Delta^* w = -\delta_o\} \quad (135)$$

and

$$\text{Harm}(G) := \{u \in \mathcal{H}_E \mid \Delta^* u = 0\}. \quad (136)$$

Here “Mono” is short for “monopole”; and “Harm” is short for “harmonic.”

Proposition 7.4. *Let (G, c_G) and (H, c_H) be weighted graphs, and let φ, ψ be a pair of functions satisfying the conditions in (127) and (128). Assume the two functions are compatible such that T in (129) is isometric $\mathcal{H}_E(H) \hookrightarrow \mathcal{H}_E(G)$, and satisfies the intertwining property (130).*

Then T maps $\text{Harm}(H)$ into $\text{Harm}(G)$, and $\text{Mono}(H)$ into $\text{Mono}(G)$.

Proof. The two assertions in the proposition are the following inclusions:

$$T(\text{Harm}(H)) \subseteq \text{Harm}(G), \quad (137)$$

and

$$T(\text{Mono}(H)) \subseteq \text{Mono}(G). \quad (138)$$

Now take $u \in \text{Harm}(H)$; i.e., $\Delta_H u = 0$. An application of (130) yields:

$$\psi(x)(\Delta_G T u)(x) = (T \Delta_H u)(x) = 0.$$

Since $\Delta_H u \equiv 0$ and $\psi(x) \neq 0$, we conclude that $\Delta_G T u = 0$, i.e., that

$$T u \in \text{Harm}(G).$$

For the second inclusion, we pick points o_H and o_G in the respective vertex sets such that $\varphi(o_G) = o_H$.

If $w \in \text{Mono}(H)$, then a second application of (130) shows that

$$\Delta_G Tw = -\psi(o_G) \delta_{o_G};$$

and so $Tw \in \text{Mono}(G)$. $\square$

Corollary 7.5. *For the dyadic tree G in Figures 2 and 3 with constant conductance, we have*

$$\text{Harm}(G) \neq 0, \text{ and } \text{Mono}(G) \neq 0.$$

Proof. A direct application of the proposition. $\square$

VIII. TWO HILBERT SPACES

Since the summations from (114) in Proposition 6.2, and on the right-hand side in Theorem 6.3, involve ℓ^2 -considerations, we now turn to a comparison of the two Hilbert spaces ℓ^2 and $\mathcal{H}_E$.

Recall the data (G, c) , Δ and $\mathcal{H}_E$ are as described above; in particular

$$c : G^1 \rightarrow \mathbb{R}_+ \quad (139)$$

is a fixed function, and the conditions listed in Definition 2.3 are assumed to hold. Note further that by Definition 2.3(c), the norm, and the inner product, $\|\cdot\|_{\mathcal{H}_E}$ and $\langle \cdot, \cdot \rangle_E$ (“E” for energy) are weighted with the fixed function c in (139); and the definitions in the energy Hilbert space $\mathcal{H}_E$ involve differences $u(x) - u(y)$ whenever $(x, y) \in G^1$.

In contrast the norm $\|\cdot\|_2$ and inner product $\langle \cdot, \cdot \rangle_{\ell^2}$ are unweighted; in fact we have:

$$\|u\|_2^2 = \sum_{x \in G^0} |u(x)|^2; \quad (140)$$

and

$$\langle u, v \rangle_{\ell^2} = \sum_{x \in G^0} \overline{u(x)} v(x), \text{ for all } u, v \in \ell^2 = \ell^2(G^0). \quad (141)$$

It is immediate that

$$\mathcal{D}_2 := \text{span} \{ \delta_x \mid x \in G^0 \} \quad (142)$$

is a dense linear subspace in ℓ^2 , and further that the Laplace operator Δ from (14) satisfies

$$\langle u, \Delta u \rangle_{\ell^2} \geq 0 \text{ for all } u \in \mathcal{D}_2. \quad (143)$$

In other words, it defines a semibounded Hermitian operator with dense domains in ℓ^2 . Moreover, by [22], this operator Δ (with domain $\mathcal{D}_2$) is essentially selfadjoint in ℓ^2 . This means that its graph closure in $\ell^2 \times \ell^2$ is a selfadjoint operator in ℓ^2 .

However, as we show in sections III–IV above, Δ is not essentially selfadjoint when viewed as an unbounded operator with dense domain in $\mathcal{H}_E$; see especially Theorem 6.3.

Our terminology for the operator Δ in equation (14) will involve some ambiguity. In fact, Δ will simultaneously be viewed as a densely defined operator in ℓ^2 , and in $\mathcal{H}_E$; the Hilbert spaces are different, and the respective dense domains are different as well. Each of the two operators will be closed; the first in the graph norm in $\ell^2 \times \ell^2$, and the second in $\mathcal{H}_E \times \mathcal{H}_E$. The first of the two closed operators is selfadjoint, while the second generally is not. What is worse, the two Hilbert spaces ℓ^2 and $\mathcal{H}_E$ do not lend themselves to a direct comparison.

We shall need the following:

Lemma 8.1.

- (a) *Let Δ_2 denote the selfadjoint operator in ℓ^2 , and let $\text{dom}(\Delta_2)$ be its dense domain. Then the sum-operator $I + \Delta_2$ is invertible, and the following hold:*

$$(I + \Delta_2)(\text{dom}(\Delta_2)) = \ell^2, \quad (144)$$

and

$$(I + \Delta_2)^{-1} \ell^2 = \text{dom}(\Delta_2). \quad (145)$$

- (b) *Let Δ_E denote the closed operator version of Δ with dense domain $\text{dom}(\Delta_E)$ in $\mathcal{H}_E$. Then the following hold:*

$$(I + \Delta_E)(\text{dom}(\Delta_E)) = \mathcal{H}_E \ominus \text{Def}, \quad (146)$$

and $(I + \Delta_E)^{-1}$ is well-defined on this closed subspace.

- (c) *The operator $(I + \Delta_2)^{-1}$ is contractive from ℓ^2 to ℓ^2 ; i.e.,*

$$\|(I + \Delta_2)^{-1} u\|_2 \leq \|u\|_2 \text{ for all } u \in \ell^2. \quad (147)$$

- (d) *The operator $(I + \Delta_E)^{-1}$ is contractive from its domain (146) into $\mathcal{H}_E$, i.e.,*

$$\|(I + \Delta_E)^{-1} u\|_E \leq \|u\|_E \quad (148)$$

holds for all u of the form $u = v + \Delta_E v$.

Proof. The verification of the conclusions listed in (a)–(d) follow by combining the properties of Δ which we derived in sections III and V–VI above. $\square$

Theorem 8.2. *Let ℓ^2 , $\mathcal{H}_E$, and Δ be as specified in the lemma where Δ will have the meaning which is dictated by the contexts of the respective Hilbert spaces.*

(a) *Then we have the following containment*

$$(I + \Delta_2)^{-1} \ell^2 = \{u \in \ell^2 \mid \Delta u \in \ell^2\} \subset \mathcal{H}_E. \quad (149)$$

(b) *If $u \in \ell^2$ and $\Delta u \in \ell^2$, then*

$$\mathcal{E}(u) = \|u\|_{\mathcal{H}_E}^2 = \sum_{x \in G^0} \overline{u(x)} (\Delta u)(x). \quad (150)$$

Proof. Let $u \in (I + \Delta_2)^{-1} \ell^2$ and pick $v \in \ell^2$ such that $u + \Delta u = v$. When the inner product of ℓ^2 is considered, this means that u is in the domain of Δ_2^* , and $\Delta_2^* u = v - u \in \ell^2$. But Δ_2 is essentially selfadjoint by the lemma, so Δ_2^* is the closure of Δ_2 . Moreover $\Delta^* u$ is given by the formula (14) in section II.

Conversely, suppose u and Δu are in ℓ^2 ; then set $v := u + \Delta u$, and note that $(I + \Delta_2)^{-1} v = u$.

Our aim is now to show that the double-summation (12) in Definition 2.3 is finite; and so $u \in \mathcal{H}_E$. The conditions on the function c are as specified in Definition 2.3(a).

When the double-summation is carried out, we note that Fubini applies; we get

$$\begin{aligned} & \frac{1}{2} \sum_{\substack{\text{all } x, y \\ \text{s.t. } x \sim y}} c(x, y) |u(x) - u(y)|^2 \\ &= \sum_x \left(c(x) |u(x)|^2 - \sum_{y \sim x} \left(c(x, y) \overline{u(x)} u(y) \right) \right) \\ &= \sum_x \overline{u(x)} \left(c(x) u(x) - \sum_{y \sim x} c(x, y) u(y) \right) \\ &= \sum_x \overline{u(x)} (\Delta u)(x). \end{aligned}$$

□

Corollary 8.3. *Let ℓ^2 , $\mathcal{H}_E$, and Δ be as specified in the theorem, and let $x \in G^0$ be given.*

(a) *Then the space of solutions $u \in \mathcal{H}_E$ to the equation*

$$\Delta u = -u \text{ in } G^0 \setminus \{x\}, \quad (151)$$

is one-dimensional. (Note that (151) is the deficiency equation in the punctured vertex set $G^0 \setminus \{x\}$.)

(b) *The solution space in $\mathcal{H}_E$ from (151) is spanned by $(I + \Delta)^{-1} \delta_x$.*

Proof. The essential point is that if

$$u_x = (I + \Delta_2)^{-1} (\delta_x) \quad (152)$$

is computed first in ℓ^2 , then the two vectors u_x and $\Delta u_x = \delta_x - u_x$ are both in ℓ^2 . As a result (see Theorem 8.2(a)) we conclude that u_x is in $\mathcal{H}_E$. It follows that

$$u_x + \Delta u_x = \delta_x \quad (153)$$

holds pointwise on G^0 , and so

$$u_x + \Delta u_x = 0 \text{ on } G^0 \setminus \{x\}.$$

Conversely, if $u \in \mathcal{H}_E$ satisfies (151) then it follows that for some constant C we have $u + \Delta u = C\delta_x$; and therefore

$$u = C(I + \Delta)^{-1} \delta_x = Cu_x. \quad (154)$$

This proves that the solution space in (151) is one-dimensional and spanned by the single function $u_x (\in \mathcal{H}_E)$ in (152). $\square$

ACKNOWLEDGMENT

The author is happy to thank his colleagues (for suggestions) in the math physics and operator theory seminars at the University of Iowa, where he benefitted from numerous helpful discussions of various aspects of the results in the paper, and their applications. And he thanks Doug Slauson for typesetting the paper.

This work supported in part by the US National Science Foundation.

REFERENCES

- [1] S. Albeverio, P. E. T. Jorgensen, and A. M. Paolucci, *Multiresolution wavelet analysis of integer scale Bessel functions*, J. Math. Phys. **48** (2007), no. 7, 073516, 24. MR MR2337697 (2008e:42060)
- [2] Daniel Alpay and David Levanony, *On the reproducing kernel Hilbert spaces associated with the fractional and bi-fractional Brownian motions*, Potential Anal. **28** (2008), no. 2, 163–184. MR MR2373103
- [3] ———, *Rational functions associated with the white noise space and related topics*, Potential Anal. **29** (2008), no. 2, 195–220. MR MR2430613
- [4] Daniel Alpay, Michael Shapiro, and Dan Volok, *Reproducing kernel spaces of series of Fueter polynomials*, Operator theory in Krein spaces and nonlinear eigenvalue problems, Oper. Theory Adv. Appl., vol. 162, Birkhäuser, Basel, 2006, pp. 19–45. MR MR2240272 (2007k:46048)
- [5] N. Aronszajn, *Theory of reproducing kernels*, Trans. Amer. Math. Soc. **68** (1950), 337–404. MR MR0051437 (14,479c)
- [6] Kendall Atkinson and Weimin Han, *Theoretical numerical analysis*, second ed., Texts in Applied Mathematics, vol. 39, Springer, New York, 2005, A functional analysis framework. MR MR2153422 (2006a:65001)

- [7] Viviane Baladi, *Positive transfer operators and decay of correlations*, Advanced Series in Nonlinear Dynamics, vol. 16, World Scientific Publishing Co. Inc., River Edge, NJ, 2000. MR MR1793194 (2001k:37035)
- [8] Martin T. Barlow, Richard F. Bass, Zhen-Qing Chen, and Moritz Kassmann, *Non-local Dirichlet forms and symmetric jump processes*, Trans. Amer. Math. Soc. **361** (2009), no. 4, 1963–1999. MR MR2465826
- [9] Sara Brofferio and Wolfgang Woess, *Green kernel estimates and the full Martin boundary for random walks on lamplighter groups and Diestel-Leader graphs*, Ann. Inst. H. Poincaré Probab. Statist. **41** (2005), no. 6, 1101–1123. MR MR2172211 (2007a:60045)
- [10] Ilwoo Cho and Palle Jorgensen, *C^* -algebras generated by partial isometries*, J. Appl. Math. Comput. **26** (2008), no. 1-2, 1–48. MR MR2383652
- [11] Nelson Dunford and Jacob T. Schwartz, *Linear operators. Part II*, Wiley Classics Library, John Wiley & Sons Inc., New York, 1988, Spectral theory. Selfadjoint operators in Hilbert space, With the assistance of William G. Bade and Robert G. Bartle, Reprint of the 1963 original, A Wiley-Interscience Publication. MR MR1009163 (90g:47001b)
- [12] Dorin Ervin Dutkay, Deguang Han, Gabriel Picioroaga, and Qiyu Sun, *Orthonormal dilations of Parseval wavelets*, Math. Ann. **341** (2008), no. 3, 483–515. MR MR2399155 (2009c:42080)
- [13] Dorin Ervin Dutkay and Palle E. T. Jorgensen, *Harmonic analysis and dynamics for affine iterated function systems*, Houston J. Math. **33** (2007), no. 3, 877–905. MR MR2335741
- [14] ———, *Martingales, endomorphisms, and covariant systems of operators in Hilbert space*, J. Operator Theory **58** (2007), no. 2, 269–310. MR MR2358531
- [15] ———, *A duality approach to representations of Baumslag-Solitar groups*, Group representations, ergodic theory, and mathematical physics: a tribute to George W. Mackey, Contemp. Math., vol. 449, Amer. Math. Soc., Providence, RI, 2008, pp. 99–127. MR MR2391800
- [16] ———, *Fourier series on fractals: a parallel with wavelet theory*, Radon transforms, geometry, and wavelets, Contemp. Math., vol. 464, Amer. Math. Soc., Providence, RI, 2008, pp. 75–101. MR MR2440130
- [17] ———, *Quasiperiodic spectra and orthogonality for iterated function system measures*, Math. Z. **261** (2009), no. 2, 373–397. MR MR2457304
- [18] Takeyuki Hida, *Brownian motion*, Applications of Mathematics, vol. 11, Springer-Verlag, New York, 1980, Translated from the Japanese by the author and T. P. Speed. MR MR562914 (81a:60089)
- [19] Palle E. T. Jorgensen, Keri Kornelson, and Karen Shuman, *Harmonic analysis of iterated function systems with overlap*, J. Math. Phys. **48** (2007), no. 8, 083511, 35. MR MR2349423
- [20] ———, *Orthogonal exponentials for Bernoulli iterated function systems*, Representations, wavelets, and frames, Appl. Numer. Harmon. Anal., Birkhäuser Boston, Boston, MA, 2008, pp. 217–237. MR MR2459320
- [21] Palle E. T. Jorgensen and Anilesh Mohari, *Localized bases in $L^2(0, 1)$ and their use in the analysis of Brownian motion*, J. Approx. Theory **151** (2008), no. 1, 20–41. MR MR2403894 (2009c:42086)
- [22] Palle E. T. Jorgensen and Erin P. J. Pearse, *Operator theory of electrical resistance networks*, arXiv:0806.3881v1 [math.OA].

- [23] Palle E. T. Jorgensen and Myung-Sin Song, *Entropy encoding, Hilbert space, and Karhunen-Loève transforms*, J. Math. Phys. **48** (2007), no. 10, 103503, 22. MR MR2362796 (2008k:94016)
- [24] A. Kolmogoroff, *Grundbegriffe der Wahrscheinlichkeitsrechnung*, Springer-Verlag, Berlin, 1977, Reprint of the 1933 original. MR MR0494348 (58 #13242)
- [25] Peter D. Lax and Ralph S. Phillips, *Scattering theory for automorphic functions*, Bull. Amer. Math. Soc. (N.S.) **2** (1980), no. 2, 261–295. MR MR555264 (81c:10037)
- [26] Edward Nelson, *The free Markoff field*, J. Functional Analysis **12** (1973), 211–227. MR MR0343816 (49 #8556)
- [27] Ronald Ortner and Wolfgang Woess, *Non-backtracking random walks and cogrowth of graphs*, Canad. J. Math. **59** (2007), no. 4, 828–844. MR MR2338235 (2008h:05057)
- [28] Michael Reed and Barry Simon, *Methods of modern mathematical physics. II. Fourier analysis, self-adjointness*, Academic Press [Harcourt Brace Jovanovich Publishers], New York, 1975. MR MR0493420 (58 #12429b)
- [29] Myung-Sin Song, *Wavelet image compression*, Operator theory, operator algebras, and applications, Contemp. Math., vol. 414, Amer. Math. Soc., Providence, RI, 2006, pp. 41–73. MR MR2270250 (2008h:42078)
- [30] ———, *Entropy encoding in wavelet image compression*, Representations, wavelets, and frames, Appl. Numer. Harmon. Anal., Birkhäuser Boston, Boston, MA, 2008, pp. 293–311. MR MR2459323
- [31] Marshall Harvey Stone, *Linear transformations in Hilbert space*, American Mathematical Society Colloquium Publications, vol. 15, American Mathematical Society, Providence, RI, 1990, Reprint of the 1932 original. MR MR1451877 (99k:47001)
- [32] J. von Neumann, *Über adjungierte Funktionaloperatoren*, Ann. of Math. (2) **33** (1932), no. 2, 294–310. MR MR1503053
- [33] Kazuhito Yamasaki and Hiroyuki Nagahama, *Energy integral in fracture mechanics (J-integral) and Gauss-Bonnet theorem*, ZAMM Z. Angew. Math. Mech. **88** (2008), no. 6, 515–520. MR MR2423340
- [34] Haizhang Zhang, *Orthogonality from disjoint support in reproducing kernel Hilbert spaces*, J. Math. Anal. Appl. **349** (2009), no. 1, 201–210. MR MR2455741

DEPARTMENT OF MATHEMATICS, 14 MLH, THE UNIVERSITY OF IOWA, IOWA CITY, IA 52242-1419, USA

E-mail address: jorgen@math.uiowa.edu

URL: <http://www.math.uiowa.edu>